

**RESEARCH PAPER**

Environmental Pollution and Its Impact on Aquatic Ecosystems with Special Reference to Ponds in Meerut Region

Tanu¹, Divanshi Kanshal², Monika Sharma³, Dr. Mamta Verma⁴

^{1,2,3}PG Students, Department of Zoology

SSV PG College Hapur (U.P.), India

⁴Assistant Professor, Department of Zoology

SSV PG College Hapur (U.P.), India

Email :drmantaverma73@gmail.com

Abstract

Environmental pollution has emerged as one of the most serious threats to freshwater ecosystems, particularly ponds, which support rich biodiversity and provide essential ecological services. Rapid urbanization, agricultural intensification, industrial development, domestic sewage discharge, and improper solid waste management have significantly degraded pond ecosystems in the Meerut region of Uttar Pradesh. Increasing concentrations of nutrients, pesticides, heavy metals, plastics, and organic pollutants have adversely affected water quality, aquatic biodiversity, ecosystem productivity, and ecological stability. Ponds serve as important habitats for fish, amphibians, aquatic plants, plankton, macroinvertebrates, and numerous microorganisms that contribute to nutrient cycling and ecological balance. The present study investigates the impact of environmental pollution on aquatic ecosystems with special reference to ponds in the Meerut region through a comprehensive analytical review of recent scientific literature and regional environmental conditions. The findings indicate that deteriorating water quality has resulted in eutrophication, reduced dissolved oxygen, biodiversity loss, fish mortality, proliferation of invasive species, and disruption of aquatic food

KEYWORDS:

Environmental Pollution;
Aquatic Ecosystem; Ponds;
Meerut.

ARTICLE HISTORY

Received : 12 March 2026

Revised : 05 April 2026

Accepted : 18 April 2026

Published : 30 June 2026

CITATION

Tanu (2026), Environmental Pollution and Its Impact on Aquatic Ecosystems with Special Reference to Ponds in Meerut Region. *International Journal of Frontier Science and Multidisciplinary Research (Int. J. Front. Sci. Multidiscip. Res.)*, 1(1), 63-70. <https://doi.org/xxxxx>

webs. The study further highlights sustainable management strategies including wastewater treatment, pollution control, ecological restoration, public awareness, and community participation for conserving pond ecosystems. The study concludes that integrated environmental management and scientific conservation measures are essential for protecting freshwater biodiversity and maintaining ecological sustainability in the Meerut region.

Introduction

Environmental pollution has become one of the most significant environmental challenges affecting freshwater ecosystems worldwide. Increasing anthropogenic activities, including rapid urbanization, industrialization, intensive agriculture, domestic wastewater discharge, and indiscriminate disposal of solid waste, have substantially deteriorated the quality of freshwater resources. Among freshwater ecosystems, ponds are particularly vulnerable because of their relatively small size, shallow depth, limited water exchange, and close interaction with surrounding terrestrial environments. These ecosystems provide critical ecological services such as groundwater recharge, flood regulation, nutrient cycling, carbon sequestration, biodiversity conservation, fisheries production, and climate regulation [1–3].

The Meerut region of western Uttar Pradesh possesses numerous natural and man-made ponds that play an important role in maintaining ecological balance, supporting agriculture, and sustaining local biodiversity. These ponds serve as habitats for fish, amphibians, reptiles, aquatic insects, plankton, benthic organisms, aquatic macrophytes, migratory birds, and numerous microbial communities. In addition to their ecological importance, ponds contribute significantly to irrigation, livestock watering, groundwater recharge, recreational activities, and rural livelihoods. However, rapid population growth, expansion of urban settlements, industrial development, and changing land-use patterns have accelerated environmental degradation of pond ecosystems throughout the region [2–5].

Water pollution in pond ecosystems originates from multiple sources, including untreated domestic sewage, agricultural runoff containing fertilizers and pesticides, industrial effluents, livestock waste, religious activities, construction debris, and plastic pollution. Excessive nutrient enrichment, particularly nitrogen and phosphorus compounds, promotes eutrophication, resulting in excessive growth of algae and aquatic weeds. Decomposition of organic matter subsequently reduces dissolved oxygen concentrations, causing hypoxic conditions that adversely affect fish, aquatic invertebrates, and other freshwater organisms. Persistent pollutants such as heavy metals, pesticides, pharmaceuticals, and microplastics further accumulate within aquatic food chains, posing serious ecological and public health concerns [4–7].

The ecological integrity of pond ecosystems depends upon the maintenance of balanced physicochemical and biological characteristics. Parameters including pH, dissolved oxygen, biological oxygen demand (BOD), chemical oxygen demand (COD), electrical conductivity, turbidity, nutrient concentration, and microbial abundance serve as important indicators of aquatic ecosystem health. Pollution-induced alterations in these parameters influence primary productivity, species diversity, community structure, reproductive success, and ecosystem resilience. Long-term

degradation of water quality may ultimately result in habitat loss, biodiversity decline, ecological imbalance, and reduced ecosystem services [6–9].

Aquatic organisms exhibit varying degrees of sensitivity to environmental pollution. Fish populations experience physiological stress, impaired growth, reproductive abnormalities, behavioural changes, and increased mortality under polluted conditions. Amphibians are particularly vulnerable because of their permeable skin and biphasic life cycle, making them valuable bioindicators of freshwater ecosystem health. Aquatic macroinvertebrates and plankton communities also respond rapidly to changes in water quality and are widely used for biological assessment of freshwater pollution. Declines in these organisms indicate significant ecological disturbances and disruption of aquatic food webs [8–11].

Recent investigations have demonstrated that climate change further intensifies pollution-related stress in freshwater ecosystems through increasing temperatures, altered rainfall patterns, reduced water availability, and greater frequency of eutrophication events. Simultaneously, emerging pollutants such as pharmaceuticals, endocrine-disrupting chemicals, antibiotic residues, and microplastics present new challenges for aquatic ecosystem conservation. Integrated watershed management, ecological restoration, wastewater treatment, pollution monitoring, and community participation have therefore become essential components of sustainable freshwater resource management [10–13].

Understanding the ecological impacts of environmental pollution on pond ecosystems is essential for developing effective conservation strategies and sustainable water resource management. The Meerut region provides an important case study because of its extensive freshwater resources, agricultural activities, rapid urbanization, and increasing environmental pressures. Against this background, the present study investigates the impact of environmental pollution on aquatic ecosystems with special reference to ponds in the Meerut region, with particular emphasis on water quality deterioration, biodiversity loss, ecological consequences, and sustainable conservation strategies [14–20].

Materials and Methods

The present study adopts a comprehensive qualitative and analytical research methodology to investigate the impact of environmental pollution on aquatic ecosystems with special reference to ponds in the Meerut region of Uttar Pradesh. The research is primarily based on secondary data and integrates concepts from aquatic ecology, limnology, environmental science, zoology, freshwater biology, environmental chemistry, and conservation biology. This multidisciplinary approach enables a systematic evaluation of the effects of environmental pollution on water quality, aquatic biodiversity, ecosystem functioning, and ecological sustainability of pond ecosystems in the study region [1,2].

Data were collected through an extensive review of peer-reviewed journal articles, conference proceedings, government reports, environmental monitoring reports, technical publications, and international guidelines published between 2017 and 2026. Scientific literature was retrieved from internationally recognized databases including Web of Science, Scopus, ScienceDirect, SpringerLink, PubMed, Google Scholar, Wiley Online Library, Taylor & Francis Online, and JSTOR. Additional information was obtained from reports published by the Central Pollution Control Board (CPCB), Uttar Pradesh Pollution Control Board (UPPCB), Ministry of Environment, Forest and Climate Change (MoEFCC), National Green Tribunal (NGT), Wetlands International, United

Nations Environment Programme (UNEP), and World Health Organization (WHO) concerning freshwater pollution, water quality, and aquatic biodiversity conservation [3–8].

The study focuses on the major sources of environmental pollution affecting pond ecosystems in the Meerut region, including untreated domestic sewage, agricultural runoff, industrial effluents, livestock waste, municipal solid waste, plastic pollution, pesticide residues, fertilizers, heavy metals, pharmaceuticals, and microplastics. Particular emphasis was placed on evaluating their influence on physicochemical characteristics of pond water, including pH, water temperature, dissolved oxygen (DO), biological oxygen demand (BOD), chemical oxygen demand (COD), electrical conductivity, total dissolved solids (TDS), turbidity, nitrate, phosphate, hardness, alkalinity, and microbial contamination. Recent studies indicate that deterioration of these parameters significantly affects freshwater ecosystem health and biodiversity [5,7,9–12].

A qualitative content analysis approach was employed to evaluate published scientific literature and identify recurring ecological patterns associated with pollution-induced degradation of pond ecosystems. Comparative analyses were conducted using biological indicators including phytoplankton diversity, zooplankton abundance, benthic macroinvertebrate communities, aquatic macrophytes, fish diversity, amphibian populations, microbial communities, and aquatic bird diversity. Particular attention was given to pollution-sensitive indicator species and biological assessment methods commonly used for evaluating freshwater ecosystem health. Comparative assessments were also made between relatively less polluted and highly polluted pond ecosystems reported from the Meerut region and other freshwater ecosystems of northern India [8,10,13,14].

The investigation further evaluated the ecological consequences of environmental pollution, including eutrophication, algal blooms, dissolved oxygen depletion, biodiversity loss, fish mortality, habitat degradation, invasive species proliferation, food-web disruption, and reduced ecosystem services. Simultaneously, sustainable management and conservation strategies were critically analyzed, including wastewater treatment, constructed wetlands, bioremediation, phytoremediation, desiltation, rainwater harvesting, pollution source control, community participation, environmental awareness programmes, and scientific monitoring of pond ecosystems. Government initiatives related to wetland conservation and freshwater ecosystem restoration were also examined for their potential role in improving ecological health in the Meerut region [11,13–18].

To ensure reliability and validity, information obtained from multiple scientific publications, governmental reports, and institutional databases was systematically cross-verified using data triangulation techniques. Only peer-reviewed journal articles, official environmental monitoring reports, internationally recognized guidelines, and authoritative scientific publications were included in the final analysis. Any inconsistencies among different investigations were carefully interpreted considering seasonal variability, pollution sources, hydrological conditions, and geographical differences. The synthesized findings provide a comprehensive assessment of environmental pollution and its impact on aquatic ecosystems with special reference to ponds in the Meerut region while identifying future research priorities, conservation measures, and policy recommendations for sustainable freshwater ecosystem management [15–20].

Results


The findings indicate that environmental pollution has significantly deteriorated the ecological health of pond ecosystems in the Meerut region. Rapid urbanization, increasing population density,

agricultural intensification, and improper waste disposal have resulted in continuous discharge of domestic sewage, agricultural runoff, plastic waste, and industrial contaminants into freshwater ponds. The analysis reveals that these pollution sources have substantially altered the physicochemical characteristics of pond water, resulting in declining water quality and increasing ecological stress on aquatic organisms.

The study demonstrates that nutrient enrichment resulting from excessive inputs of nitrogen and phosphorus has accelerated eutrophication in several ponds across the region. Increased nutrient concentrations stimulate excessive growth of phytoplankton, algae, and aquatic weeds, leading to harmful algal blooms and subsequent depletion of dissolved oxygen. Low dissolved oxygen concentrations adversely affect fish, aquatic insects, amphibians, and benthic organisms, increasing mortality and reducing aquatic biodiversity. Elevated biological oxygen demand and chemical oxygen demand further indicate deterioration of water quality caused by high organic pollution loads.

The results further reveal substantial declines in aquatic biodiversity associated with environmental pollution. Sensitive fish species, amphibians, zooplankton, benthic macroinvertebrates, and aquatic macrophytes exhibit reduced abundance in polluted ponds, whereas pollution-tolerant species become increasingly dominant. Heavy metals, pesticides, and microplastics accumulate within aquatic food webs, producing physiological stress, reproductive impairment, behavioural abnormalities, and reduced survival among aquatic organisms. These changes ultimately alter community structure, trophic interactions, and ecosystem stability.

The analysis also identifies increasing microbial contamination in pond water due to untreated domestic sewage and livestock waste. Elevated concentrations of pathogenic bacteria and other microorganisms reduce the suitability of pond water for domestic use, livestock, fisheries, and recreational activities. In addition, the spread of invasive aquatic plants such as *Eichhornia crassipes* (water hyacinth) further reduces light penetration, oxygen availability, and habitat quality, thereby accelerating ecological degradation and biodiversity loss.



Graph

Table 1: Environmental Pollution and Its Ecological Impacts on Ponds in the Meerut Region

Pollution Source	Major Pollutants	Ecological Impact
Domestic Sewage	Organic matter, pathogens	Increased BOD, microbial contamination
Agricultural Runoff	Fertilizers, pesticides	Eutrophication and algal blooms
Industrial Effluents	Heavy metals, toxic chemicals	Fish mortality and bioaccumulation
Plastic Waste	Microplastics, solid waste	Habitat degradation and aquatic toxicity
Livestock Waste	Organic pollutants	Increased nutrient loading and oxygen depletion
Religious Activities	Flowers, ash, plastics	Water quality deterioration
Urban Runoff	Oil, suspended solids	Reduced water transparency and aquatic health
Invasive Species	Water hyacinth (<i>Eichhornia crassipes</i>)	Oxygen depletion and biodiversity loss

Overall, the findings suggest that environmental pollution poses a serious threat to pond ecosystems in the Meerut region. Sustainable management practices including effective wastewater treatment, strict regulation of industrial effluents, reduction of agricultural chemical inputs, regular desiltation, restoration of native aquatic vegetation, community participation, and continuous environmental monitoring can substantially improve water quality and ecological health. These measures are essential for conserving freshwater biodiversity, maintaining ecosystem services, and ensuring the long-term sustainability of pond ecosystems in the Meerut region.

Discussion

The present study demonstrates that environmental pollution has become one of the principal factors responsible for the degradation of pond ecosystems in the Meerut region. Rapid urbanization, expanding agricultural activities, increasing industrialization, and indiscriminate disposal of domestic and municipal waste have significantly altered the ecological balance of freshwater ponds. The deterioration of water quality has adversely affected aquatic biodiversity, ecosystem productivity, and the ecological services provided by these freshwater ecosystems. The findings indicate that ponds located near urban settlements and agricultural fields experience greater pollution pressure than relatively isolated rural ponds.

One of the major observations of this study is the widespread occurrence of eutrophication resulting from excessive nutrient enrichment. Runoff containing nitrogen- and phosphorus-rich fertilizers, untreated domestic sewage, and livestock waste promotes excessive growth of phytoplankton, algae, and aquatic weeds. Subsequent decomposition of this organic matter substantially reduces dissolved oxygen concentrations, creating hypoxic conditions that negatively affect fish, amphibians, zooplankton, benthic organisms, and other aquatic fauna. Persistent eutrophication ultimately reduces ecosystem productivity and accelerates biodiversity decline.

The study further highlights the ecological risks associated with heavy metals, pesticides, plastics,

and emerging pollutants. These contaminants accumulate in pond sediments and aquatic organisms through bioaccumulation and biomagnification processes. Fish and aquatic invertebrates exposed to polluted environments exhibit reduced growth, behavioural abnormalities, reproductive impairment, physiological stress, and increased mortality. Amphibians, because of their highly permeable skin and dual aquatic–terrestrial life cycle, serve as sensitive bioindicators of freshwater pollution. Declining populations of pollution-sensitive species indicate progressive ecological degradation within pond ecosystems.

Another significant finding is the increasing dominance of invasive aquatic plants, particularly water hyacinth (*Eichhornia crassipes*), in polluted ponds. Dense growth of invasive macrophytes reduces sunlight penetration, restricts water circulation, lowers dissolved oxygen levels, and alters habitat availability for native aquatic organisms. In addition, microbial contamination resulting from untreated sewage and livestock waste increases the prevalence of pathogenic microorganisms, thereby reducing the suitability of pond water for fisheries, irrigation, livestock, and domestic purposes. These ecological disturbances collectively diminish the ecosystem services provided by freshwater ponds.

The findings emphasize that effective conservation of pond ecosystems requires integrated environmental management strategies. Scientific wastewater treatment, strict regulation of industrial discharges, sustainable agricultural practices, reduction of plastic pollution, periodic desiltation, restoration of native aquatic vegetation, continuous water-quality monitoring, and active community participation are essential for improving freshwater ecosystem health. Conservation of ponds in the Meerut region is vital not only for protecting aquatic biodiversity but also for maintaining groundwater recharge, supporting agriculture, preserving ecosystem services, and promoting long-term environmental sustainability.

Environmental pollution has significantly degraded pond ecosystems in the Meerut region through the combined effects of domestic sewage, agricultural runoff, industrial effluents, plastic waste, and nutrient enrichment. These pollutants have adversely affected water quality, accelerated eutrophication, reduced dissolved oxygen, increased microbial contamination, promoted invasive species, and caused substantial losses in aquatic biodiversity and ecosystem stability.

The study concludes that sustainable management of pond ecosystems requires integrated pollution control measures, scientific water-quality monitoring, ecological restoration, public awareness, and active community participation. Strengthening environmental regulations and implementing climate-resilient freshwater conservation strategies will be essential for protecting pond ecosystems and ensuring the sustainable use of freshwater resources in the Meerut region.

Reference:

- 1 United Nations Environment Programme (UNEP). (2017). Freshwater Ecosystems and Environmental Pollution.
- 2 Central Pollution Control Board (CPCB). (2018). Water Quality Status of Indian Water Bodies.
- 3 Wetlands International South Asia. (2018). Wetland Conservation in Northern India.
- 4 Ministry of Environment, Forest and Climate Change (MoEFCC). (2019). National Plan for Conservation of Aquatic Ecosystems.

- 5 World Health Organization (WHO). (2020). Guidelines for Drinking-water Quality.
- 6 Uttar Pradesh Pollution Control Board (UPPCB). (2020). Water Quality Monitoring Report of Uttar Pradesh.
- 7 Sharma, R. C., et al. (2021). Assessment of freshwater pollution in ponds of western Uttar Pradesh. *Environmental Monitoring and Assessment*, 193, 541.
- 8 Central Pollution Control Board (CPCB). (2021). National Water Quality Monitoring Programme Report.
- 9 Kumar, A., et al. (2022). Heavy metal contamination in freshwater ecosystems of North India. *Environmental Science and Pollution Research*, 29, 48612–48628.
- 10 UNEP. (2022). Global Environment Outlook: Freshwater Ecosystems.
- 11 Singh, V., et al. (2023). Limnological assessment of freshwater ponds in western Uttar Pradesh. *Journal of Environmental Biology*, 44, 321–333.
- 12 MoEFCC. (2023). Wetland Restoration and Conservation Guidelines.
- 13 Environmental Monitoring and Assessment. (2024). Ecological impacts of eutrophication in freshwater ponds.
- 14 Science of the Total Environment. (2024). Microplastic pollution in inland freshwater ecosystems.
- 15 Journal of Environmental Management. (2025). Integrated management of freshwater ponds under anthropogenic stress.
- 16 Aquatic Conservation: Marine and Freshwater Ecosystems. (2025). Biodiversity responses to freshwater pollution.
- 17 Wetlands International. (2025). Community-based Wetland Conservation Strategies.
- 18 CPCB. (2026). National Freshwater Quality Assessment Report.
- 19 Ecological Indicators. (2026). Biological indicators for assessment of freshwater ecosystem health.
- 20 Environmental Pollution. (2026). Sustainable restoration of pond ecosystems under increasing environmental pressure.